

## All Landscape Installation & Maintenance Professionals


GENERAL DIRECTIONS FOR USE: Apply using a rotary spreader at a rate of 1 lb. pound per 600 square feet or apply by hand with scoop in beds, mulch or rough grass. Early Sept/Early Oct apply a 20 foot band to the perimeter of the property where voles are nesting during the summer months. This helps reduce voles from entering the property. Directions for Northern Snowfall Regions: 1st application (Late Oct. after irrigation is off) apply to areas that have repeated winter vole infestation each year. 2nd application (Early Nov. - 14 days after 1st application) - Re-apply to the same areas prior to snowfall. (Vole problems seem to be heaviest in areas where snow drifting occurs). 3rd application (Early Spring/Late Winter) apply immediately after snow melt. Directions for Midwest & Southern Regions: Apply 2 applications, 2 weeks apart to areas that are showing vole damage. Repeat again 30 days after second application. SEE FULL APPLICATION INSTRUCTIONS INSIDE.

PRECAUTIONARY STATEMENTS: Wash skin after use. Avoid contact with eyes or cuts. If in eyes, flush with water 15-20 mins. Call Poison Center or doctor for treatment. Storage: Store in original sealed container. Store away from direct sunlight or excessive heat. Do not contaminate water by storage or disposal. Do not use on food or feed crops. Do not apply directly to water. Disposal: Do not reuse empty container. Dispose of properly. Buyer assumes all responsibility for safety and use not in accordance with directions. EPIC represents that this product qualifies for exemption from EPA registration under Section 25(b) of FIFRA and has not been registered by the EPA.









Drie Ca Th Clo Ros Pe Ga Pea TO

ACTIVE INGREDIENTS	
Dried Blood	14.00%
Castor Oil	6.50%
Thyme Oil	1.50%
Clove Oil	1.00%
Rosemary Oil	1.00%
Peppermint Oil	0.50%
Garlic Oil	0.10%
OTHER	
Peanut Hulls, Fish Oil	75.40%
TOTAL	
INGREDIENTS	100.0%





